



wherein X^2 is a direct bond or is selected from O, S, SO, SO₂, N(R⁹), CO, CH(OR⁹), CON(R⁹), N(R⁹)CO, N(R⁹)CON(R⁹), SO₂N(R⁹), N(R⁹)SO₂, C(R⁹)₂O, C(R⁹)₂S and N(R⁹)C(R⁹)₂, wherein R⁹ is hydrogen or (1-6C)alkyl, and Q² is aryl, aryl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl which optionally bears 1, 2 or 3 substituents, which may be the same or different, selected from trifluoromethyl, trifluoromethoxy, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, and from a group of the formula:



wherein X^3 is a direct bond or is selected from O and N(R¹¹), wherein R¹¹ is hydrogen or (1-6C)alkyl, and R¹⁰ is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl or di-[(1-6C)alkyl]amino-(1-6C)alkyl, and any heterocyclyl group within Q² optionally bears 1 or 2 oxo or thioxo substituents;

R is selected from hydrogen, amino, hydroxy, halogeno, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, carboxy, (1-6C)alkoxycarbonyl and *N*-(heterocyclyl(3-8C)cycloalkyl)carbamoyl;

R¹ is selected from hydrogen, halogeno, trifluoromethyl, trifluoromethoxy, cyano, nitro, hydroxy, amino, mercapto, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, (3-6C)alkenoylamino, N-(1-6C)alkyl-(3-6C)alkenoylamino, (3-6C)alkynoylamino, N-(1-6C)alkyl-(3-6C)alkynoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino;

R² is selected from hydrogen, halogeno, amino, hydroxy, halogeno, (1-6C)alkyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, aryl(1-6C)alkylamino, arylamino, heterocyclyl and (2-6C)alkanoylamino;

R³ is selected from hydrogen, (1-6C)alkyl, hydroxy(1-6C)alkyl, carboxy, (1-6C)alkoxycarbonyl, carbamoyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl and *N*-(heterocyclyl(3-8C)cycloalkyl)carbamoyl;

R⁵ is, independently, as defined for **R⁴** and **R⁶**, provided that **R⁵** is not halogeno;

R⁴ and **R⁶** which may be the same or different, are selected from hydrogen, halogeno, trifluoromethyl, trifluoromethoxy, cyano, isocyano, nitro, hydroxy, mercapto, amino, formyl, carboxy, carbamoyl, sulfamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, (3-6C)alkenoylamino, N-(1-6C)alkyl-(3-6C)alkenoylamino, (3-6C)alkynoylamino, N-(1-6C)alkyl-(3-6C)alkynoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, and from a group of the formula :



wherein **X⁵** is a direct bond or is selected from O, S, SO, SO₂, N(R¹²), CO, CH(OR¹²), CON(R¹²), N(R¹²)CO, SO₂N(R¹²), N(R¹²)SO₂, OC(R¹²)₂, SC(R¹²)₂ and N(R¹²)C(R¹²)₂, wherein **R¹²** is hydrogen or (1-6C)alkyl, and **Q⁴** is aryl, aryl-(1-6C)alkyl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein adjacent carbon atoms in any (2-6C)alkylene chain within an **R⁴**, **R⁵** or **R⁶** substituent are optionally separated by the insertion into the chain of a group selected from O, S, SO, SO₂, N(R¹³), CO, CH(OR¹³), CON(R¹³), N(R¹³)CO, SO₂N(R¹³), N(R¹³)SO₂, CH=CH and C≡C wherein **R¹³** is hydrogen or (1-6C)alkyl,

and wherein any CH₂=CH- or HC≡C- group within an **R⁴**, **R⁵** or **R⁶** substituent optionally bears at the terminal CH₂= or HC≡ position a substituent selected from halogeno, carboxy, carbamoyl, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl and di-[(1-6C)alkyl]amino-(1-6C)alkyl and from a group of the formula :



wherein X^6 is a direct bond or is selected from CO and $N(R^{14})CO$, wherein R^{14} is hydrogen or (1-6C)alkyl, and Q^5 is aryl, aryl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein any CH_2 or CH_3 group within a R^4 , R^5 or R^6 substituent optionally bears on each said CH_2 or CH_3 group one or more halogeno or (1-6C)alkyl substituents or a substituent selected from hydroxy, cyano, amino, carboxy, carbamoyl, (1-6C)alkoxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino or N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, or from a group of the formula :



wherein X^7 is a direct bond or is selected from O, S, SO, SO_2 , $N(R^{15})$, CO, $CH(OR^{15})$, $CON(R^{15})$, $N(R^{15})CO$, $SO_2N(R^{15})$, $N(R^{15})SO_2$, $C(R^{15})_2O$, $C(R^{15})_2S$ and $N(R^{15})C(R^{15})_2$, wherein R^{15} is hydrogen or (1-6C)alkyl, and Q^6 is aryl, aryl-(1-6C)alkyl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl,

and wherein any aryl, heteroaryl, heterocyclyl, cycloalkyl or cycloalkenyl group within a substituent on R^4 , R^5 or R^6 optionally bears 1 or more substituents, which may be the same or different, selected from halogeno, trifluoromethyl, trifluoromethoxy, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyoxy, (1-6C)alkylthio, (1-6C)alkylsulfanyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino, N-(1-6C)alkyl-(1-6C)alkanesulfonylamino, from a group of the formula :



wherein X^8 is a direct bond or is selected from O and $N(R^{17})$, wherein R^{17} is hydrogen or (1-6C)alkyl, and R^{16} is halogeno-(1-6C)alkyl, hydroxy-(1-6C)alkyl, (1-6C)alkoxy-(1-6C)alkyl, cyano-(1-6C)alkyl, amino-(1-6C)alkyl, (1-6C)alkylamino-(1-6C)alkyl, di-[(1-6C)alkyl]amino-(1-6C)alkyl, (2-6C)alkanoylamino-(1-6C)alkyl or (1-6C)alkoxycarbonylamino-(1-6C)alkyl, and from a group of the formula :



wherein X^9 is a direct bond or is selected from O, S, SO, SO_2 , $N(R^{18})$, CO, $CH(OR^{18})$, $CON(R^{18})$, $N(R^{18})CO$, $SO_2N(R^{18})$, $N(R^{18})SO_2$, $C(R^{18})_2O$, $C(R^{18})_2S$ or $N(R^{18})C(R^{18})_2$, wherein R^{18} is hydrogen or (1-6C)alkyl, and Q^7 is aryl, aryl-(1-6C)alkyl, (3-7C)cycloalkyl, (3-7C)cycloalkyl-(1-6C)alkyl, (3-7C)cycloalkenyl, (3-7C)cycloalkenyl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl or heterocyclyl-(1-6C)alkyl which optionally bears 1 or 2 substituents, which may be the same or different, selected from halogeno, trifluoromethyl, trifluoromethoxy, cyano, nitro, hydroxy, amino, carboxy, carbamoyl, (1-6C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (1-6C)alkoxy, (2-6C)alkenyloxy, (2-6C)alkynyloxy, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, (1-6C)alkoxycarbonyl, N-(1-6C)alkylcarbamoyl, N,N-di-[(1-6C)alkyl]carbamoyl, (2-6C)alkanoyl, (2-6C)alkanoyloxy, (2-6C)alkanoylamino, N-(1-6C)alkyl-(2-6C)alkanoylamino, N-(1-6C)alkylsulfamoyl, N,N-di-[(1-6C)alkyl]sulfamoyl, (1-6C)alkanesulfonylamino and N-(1-6C)alkyl-(1-6C)alkanesulfonylamino,

or when G is NR^5 , R^4 and R^5 together with the atoms to which they are attached form a fused 5- or 6- membered heteroaryl or heterocyclyl ring, and wherein said fused 5- or 6-membered ring optionally bears one or more substituents as defined for R^4 ,

and any fused 5- or 6- membered heterocyclyl ring so formed optionally bears 1 or 2 oxo or thioxo substituents,

and wherein any heterocyclyl group within any R^4 , R^5 or R^6 substituent optionally bears 1 or 2 oxo or thioxo substituents;
or a pharmaceutically-acceptable salt thereof;
provided the compound is not 4-[-2-(6-phenylimidazo[2,1-b][1,3-thiazol-5-yl)ethenyl]-2-pyrimidinamine.


2. (original) A pharmaceutical composition which comprises a compound of the Formula I, or a pharmaceutically acceptable salt thereof, as defined in claim 1 in association with a pharmaceutically-acceptable diluent or carrier.

3. (Previously canceled)

4. (Previously canceled)

5. (Previously canceled)

6. (Previously presented) A compound according to Claim 1 wherein **R** is selected from hydrogen, halogeno, carboxy, (1-6C)alkoxycarbonyl and *N*-(heterocyclyl(3-8C)cycloalkyl)carbamoyl or a pharmaceutically acceptable salt thereof.
7. (Previously presented) A compound according to Claim 1 wherein **R**¹ is selected from hydrogen, amino and (1-6C)alkyl or a pharmaceutically acceptable salt thereof.
8. (Previously presented) A compound according to Claim 1 wherein **R**² is selected from hydrogen, halogeno, hydroxy, amino, (1-6C)alkylthio, (1-6C)alkylamino, di-[(1-6C)alkyl]amino, aryl(1-6C)alkylamino, arylamino, heterocyclyl and (2-6C)alkanoylamino or a pharmaceutically acceptable salt thereof.
9. (Previously presented) A compound according to Claim 1 wherein **R**³ is selected from hydrogen, carboxy, (1-6C)alkoxycarbonyl, hydroxy(1-6C)alkyl, N-(1-6C)alkylcarbamoyl and N-(heterocyclyl(3-8C)cycloalkyl)carbamoyl or a pharmaceutically acceptable salt thereof.
10. (Previously presented) A compound according to Claim 1 wherein **R**⁴ is hydrogen and **R**⁵ is selected from (1-6C)alkyl, aryl(1-6C)alkyl, carboxy(1-6C)alkyl, heterocyclyl(1-6C)alkyl and amino(1-6C)alkyl wherein the amino group is optionally substituted by one or more (1-6C)alkyl or a pharmaceutically acceptable salt thereof.
11. (Previously canceled)
12. (Previously canceled)
13. (New) A compound according to Claim 1 as represented by Formula II




Formula II

wherein

Y, Q¹, R, R¹, R² and R³ are as defined in Claim 1;

or a pharmaceutically-acceptable salt thereof.

14. (New) A compound according to Claim 1 as represented by Formula III



Formula III

wherein

Y, Q¹, L, R, R¹, R², R³, R⁴, R⁵, r and s are as defined in Claim 1;

or a pharmaceutically-acceptable salt thereof.